

## Nodal and Hedgehog Signaling in Midline and Left-Right Axis Development

**Alexander F. Schier**, Skirball Institute, NYU School of Medicine, 540 First Avenue, NY 10016, USA

Genetic screens in zebrafish have identified two classes of genes that lead to defects resembling human holoprosencephaly (Schier, 1997): mutations leading to cyclopia (cyclopic class) and mutations affecting the ventral midline of the neural tube and the crossing of retinal axons in the diencephalon (midline/retinotectal class). Cyclopic mutants such as cyclops, one-eyed pinhead and squint have ventral forebrain deficiencies, fused eyes or a single median eye. Molecular studies have indicated that these mutations affect signaling by TGF-beta factors of the Nodal family (Schier and Shen, 2000). Cyclops and squint encode Nodal-related proteins and one-eyed pinhead affects an EGF-CFC protein (Feldman et al., 1998; Rebagliati et al., 1998; Sampath et al., 1998; Zhang et al., 1998; Gritsman et al, 1999). Midline/retinotectal mutants such as sonic you, you-too and detour affect subregions in the ventral diencephalon. Molecular studies have indicated that these three genes encode components of the hedgehog signaling pathway: sonic you disrupts the signaling molecule sonic hedgehog and you-too and detour affect the transcription factors gli2 and gli1, respectively (Schauerte et al., 1998; Karlstrom et al., 1999; Karlstrom et al., in preparation). I will discuss how Nodal and Hedgehog signaling pattern vertebrate embryos and discuss the implications for understanding holoprosencephaly.

Feldman, B., Gates, M. A., Egan, E. S., Dougan, S. T., Rennebeck, G., Sirotnik, H. I., Schier, A. F., and Talbot, W. S. (1998). Zebrafish organizer development and germ-layer formation require nodal-related signals. *Nature* 395, 181-5.

Gritsman, K., Talbot, W. S., and Schier, A. F. (2000). Nodal signaling patterns the organizer. *Development* 127, 921-932.

Gritsman, K., Zhang, J., Cheng, S., Heckscher, E., Talbot, W. S., and Schier, A. F. (1999). The EGF-CFC protein one-eyed pinhead is essential for nodal signaling. *Cell* 97, 121-132.

Karlstrom, R. O., Talbot, W. S., and Schier, A. F. (1999). Comparative synteny cloning of zebrafish you-too: mutations in the Hedgehog target gli2 affect ventral forebrain patterning. *Genes & Development* 13, 388-93.

Rebagliati, M. R., Toyama, R., Haffter, P., and Dawid, I. B. (1998). cyclops encodes a nodal-related factor involved in midline signaling. *Proc Natl Acad Sci U S A* 95, 9932-7.

Sampath, K., Rubinstein, A. L., Cheng, A. M., Liang, J. O., Fekany, K., Solnica-Krezel, L., Korzh, V., Halpern, M. E., and Wright, C. V. (1998). Induction of the zebrafish ventral brain and floorplate requires cyclops/nodal signalling. *Nature* 395, 185-9.

Schauerte, H. E., van Eeden, F. J., Fricke, C., Odenthal, J., Strähle, U., and Haffter, P. (1998). Sonic hedgehog is not required for the induction of medial floor plate cells in the zebrafish. *Development* 125, 2983-93.

Schier, A. F. (1997). Genetics of neural development in zebrafish. *Current Opinion in Neurobiology* 7, 119-26.

Schier, A. F., and Shen, M. M. (2000). Nodal signalling in vertebrate development. *Nature* 403, 385-389.

Zhang, J., Talbot, W. S., and Schier, A. F. (1998). Positional cloning identifies zebrafish one-eyed pinhead as a permissive EGF-related ligand required during gastrulation. *Cell* 92, 241-51.

Reviews:

Schier, A.F. and Shen, M.M. (2000). Nodal signalling in vertebrate development. *Nature* 403, 385-389.

Burdine, R.D. and Schier, A.F. (2000). Conserved and divergent mechanisms in left-right axis formation. *Genes & Development* 14, 763-776

Schier, A.F. (2001). Zebrafish axis formation and patterning. *Curr. Opin. Genetics and Development* 11, 393-404.

Original Articles:

Karlstrom, R.O., Talbot, W.S., and Schier, A.F. (1999). Comparative synteny cloning of zebrafish you-too: mutations in the hedgehog target gli2 affect ventral forebrain patterning. *Genes & Development* 13, 388-393.

Gritsman, K., Talbot, W.S. and Schier, A.F. (2000). Nodal signaling patterns the organizer. *Development* 127, 921-932.

Chen, Y. and Schier, A.F. (2001). The zebrafish nodal signal squint functions as a morphogen. *Nature* 411, 607-610.